

CONSIDERATIONS REGARDING GLOBAL WARMING AND CLIMATE CHANGE

Ioana Alina Crețan¹

Abstract: The paper presents global considerations regarding the global warming and climate change. It also describes the green-house gases and gas concentrations that have increased over the industrial period. In the last part it tries to predict some implication due to the climate change in all the environment factors implied

Keywords: environment factors, global warming, climate change.

1. INTRODUCTION

The terms global warming and climate change are often used interchangeably, but the two phenomena are different. Global warming is the rise in global temperatures due to an increase of heat-trapping carbon emissions in the atmosphere.

Climate change, on the other hand, is a more general term that refers to changes in many climatic factors (such as temperature and precipitation) around the world. These changes are happening at different rates and in different ways. For example, the United States has become wetter over the 20th century, while the Sahel region of central Africa has become drier.

Infrared (IR) active gases, principally water vapor (H_2O), carbon dioxide (CO_2), and ozone (O_3), naturally present in the Earth's atmosphere, absorb thermal IR radiation emitted by the Earth's surface and atmosphere. The atmosphere is warmed by this mechanism and, in turn, emits IR radiation, with a significant portion of this energy acting to warm the surface and the lower atmosphere. As a consequence the average surface air temperature of the Earth is about $30^\circ C$ higher than it would be without atmospheric absorption and reradiating of IR energy [1].

The principal greenhouse gas concentrations that have increased over the industrial period are carbon dioxide (CO_2), methane (CH_4), nitrous oxide (N_2O), and chlorofluorocarbons CFC-11 (CCl_3F) and CFC-12 (CCl_2F_2) [2]. The observed increase of CO_2 in the atmosphere from about 280 ppm in the preindustrial era to about 364 ppm in 1997 (Figure 1) [3, 4] has come largely from fossil fuel combustion and cement production. For some greenhouse gases persistence can be estimated from "mean residence times," which are obtained with simple linear models and represent the time that would be required for removal of 63% of

the anthropogenic excess of the material in the atmosphere, if anthropogenic sources were abruptly diminished to zero [5,6].




Figure 1. Concentrations of principal anthropogenic greenhouse gases in the industrial era [3,4].

This approach yields a rough measure of the persistence in the atmosphere of anthropogenic additions of CH_4 with an estimated mean residence time of 10 years [7, 8]; N_2O , 100 years and CFC-11 and CFC-12, 50 and 102 years.

Of the several anthropogenic greenhouse gases, CO_2 is the most important agent of potential future climate warming because of its large current greenhouse forcing [9], and its substantial projected future forcing [9], and its long persistence in the atmosphere (see above). Understanding climate response to a specified forcing is one of the major challenges facing the climate research community. The equilibrium response of the nonlinear climate system depends in complex ways on various feedbacks, such as changes in water vapour concentration and cloudiness that can augment or diminish climate response from that which would occur in the absence of such feedbacks.

2. CLIMATE CHANGE AND CARBON DIOXIDE.

The most commonly considered indicator of climate change is the surface air temperature.

¹ Hydrotechnical Engineering Department, "Politehnica" University of Timișoara, Romania,

Extensive efforts have been made to examine the trends in global and regional mean temperatures over time [10; 11; 12] and in the global patterns of temperature change [13, 14].

Worldwide temperature measurements, carefully screened with tools used to establish the effects of urbanization, have been used to estimate that global mean annual surface temperatures have increased between 0,3 and 0,6° C during the last 150 years [15; 16]. However, it must be stressed that the increase has not been monotonic, with annual fluctuations in the global annual mean temperature equal to an appreciable fraction of the overall rise over this time period. No single explanation can account for this variability.

Although temperature is usually the first variable considered in assessments of global climate change, it is important to consider other data that integrate the state of the climate system over space and time. These include temperature proxy data (such as tree ring records), borehole temperature measurements in soil, permafrost, and ice sheets, and measurements of the mass balance of valley glaciers and ice caps. Several recent proxy temperature reconstructions have suggested that the warming during the twentieth century is greater than any seen in the last 400 to 600 years [17] and perhaps the last 1200 to 1500

Glaciers are present on every continent except Australia (Figure 2); they are thus excellent geographically dispersed regional indicators of climate change. The Earth's valley glaciers, ice caps, and ice fields and their associated outlet glaciers have generally been shrinking and receding during the last century. Studies in North America [18], South America [19], Europe [20], Iceland [21], Africa [22], and Asia have shown substantial recession of many of the ice caps and no tidewater, no surge-type glaciers [23] since the early nineteenth century.




Figure 2. Glacial Change

The record of the past few thousand years is more difficult to piece together than the more recent record because fewer data are available. There is evidence from this period that climatic conditions were sometimes warmer and sometimes cooler than at present [24,25].

It is clear from these records, and from many other studies of paleoclimate evidence throughout the geologic record, that the global climate system has been influenced by many factors in addition to greenhouse gases [6]. To evaluate geologic evidence for the influence of greenhouse gases, one must focus

on records from periods when changes in atmospheric CO₂ were much larger than those that occurred during the millennia immediately preceding the recent increase in anthropogenic CO₂ production.

Larger natural variations in atmospheric CO₂ have been inferred from the geologic record of the more distant past [26]. Variations of 80-100 ppm, observed in analyses of gas bubbles trapped in glacier-ice cores, are correlated with the glacial (“ice age”) and interglacial climatic oscillations of the latest Pleistocene and Holocene Epochs (Figure 3); [27]. Glacial periods are associated with low CO₂ concentrations, and interglacial periods with high CO₂ concentrations. Ice core methane profiles show a similar correlation with climate [28].




Figure 3. Carbon dioxide concentration (top), proxy temperature (middle), and methane concentration from analyses of ice cores from Vostok, Antarctica [29].

Still larger past variations in atmospheric CO₂, including increases to concentrations several times higher than recent levels, have been estimated using geochemical models constrained by the sediment record [30]. During the last several hundred million years, these larger and slower CO₂ changes can be correlated with general features of climate change [30,31].

Paleoclimate model simulations (using models similar in many ways to the models used in modern climate projections) support the importance of CO₂ in explaining global mean temperatures in the geologic past [30, 32]. Model simulations have also shown the importance of changes in other climate controls, for example the configuration of Earth's orbit [25] and the geographical distribution and elevation of continental areas [33].

Significant gaps remain in understanding the relationships among these diverse climatic influences. However, the prevailing paradigm in paleoclimate research treats the radiative effects of atmospheric CO₂ as an integral component in a complex system of many variables and interactive influences on global climate.

The complexity of the long-term coupling of CO₂ and climate is enhanced by the extent to which climate variability is hypothesized to have influenced past atmospheric CO₂ concentrations.

Glacial/interglacial CO₂ variations during the Pleistocene epoch appear to have involved a combination of changes in global carbon cycling that were probably driven by some aspect of climate change [31, 34; 35, 36]. Likewise, for timescales of millions of years and longer, atmospheric CO₂ appears to have been affected by the influence of climate on weathering and erosion rates.

Thus current interpretation of the geologic record suggests that greenhouse gases both respond and contribute to climate change. Atmospheric CO₂ is viewed as one of many components of the climate system that interact in complex ways over a wide range of timescales. A change in one of these interactive components is likely to affect other aspects of the global climate system. This interactive relationship between CO₂ and climate implies that the geologic record is not likely to reveal analogs of simple climate forcing by anthropogenic CO₂ emissions [25, 36]. There is no known geologic precedent for large increases of atmospheric CO₂ without simultaneous changes in other components of the carbon cycle and climate system.

3. PREDICTIONS REGARDING CLIMATE CHANGE FROM INCREASED GREENHOUSE GASES

Some of the predicted responses to increases in greenhouse gases include increases in mean surface air temperature, increases in global mean rates of precipitation and evaporation, rising sea level, and changes in the biosphere (Figure 4). Many of these predictions are based largely on computer models that simulate fundamental geophysical processes.

Most model simulations of Earth's climate indicate that an increase in the atmospheric concentration of a greenhouse gas will lead to an increase in the average surface air temperature of the Earth [37]. For example, the 18 model runs (using 7 independent models) quoted by Kattenberg et al. predict an equilibrium temperature increase of $2.0 \pm 0.6^\circ \text{C}$ for simulations using double the current level of atmospheric CO₂.




Figure 4. More pest invading due to warmer temperatures.

An increase in surface air temperature would cause an increase in evaporation and generally higher levels of atmospheric water vapor (Figure 5). The positive feedback associated with this leads to the expectation that an increase in surface air temperatures would lead to a more intense hydrological cycle, with more frequent heavy

precipitation events [37].



Figure 5. No more snow

However, because of the coarse spatial resolution of present general circulation models, simulations of the regional and seasonal distribution of precipitation are poor.

Another possible consequence of greenhouse-gas-induced climate change is elevated sea level (Figure 6,7).



Figure 6. Vanishing Islands

The main factors that contribute to sea level rise are thermal expansion of ocean water and the melting of glaciers, both of which are in response to higher air temperatures. Although it has been well established that meltwater from the world's small glaciers has contributed to sea level rise during the last century [23], the mass balance of the ice sheets in Greenland and Antarctica is unknown.

However, recent geodetic airborne laser altimeter measurements indicate that between 1993 and 1998 the south-eastern part of the Greenland ice sheet thinned overall, with a thickening at a rate of $0.5 \pm 0.7 \text{ cm/yr}$ at elevations above 2000 m (not corrected for crustal motion) and a thinning at the low elevations at rates up to 1 m/yr [38].



Figure 7. Rising tides

Worldwide measurements from tidal gauges during the last 100 years indicate that mean sea level has risen between 10 and 25 cm (18 cm mean) [39, 40, 41]. This rate is greater than would be expected from the archaeological and geological record of sea level from the last two millennia [40]. Most modelling studies, including simulations of the combined effects of increasing greenhouse gases and aerosols, predict that the trend in rising sea level will continue in the future [40,41]. A possible biological effect may be seen in evidence that there has been an increase in the active growing season at high latitudes in the Northern Hemisphere.

REFERENCES

[1] Peixoto, J. P., and A. H. Oort, *Physics of Climate*, 520 pp., American Institute of Physics, New York, 1992.

[2] Hansen, J. E., M. Sato, A. Lacis, R. Ruedy, I. Gegen, and E. Matthews, *Climate forcings in the industrial era*, *Proceedings of the National Academy of Sciences*, 95, 12,753-12,758, 1998.

[3] Friedli, H., H. Lotscher, H. Oeschger, U. Siegenthaler, and B. Stauffer, *Ice core record of 13C/12C ratio of atmospheric CO2 in the past two centuries*, *Nature*, 324, 237-238, 1986.

[4] Keeling, C. D., and T. P. Whorf, *Atmospheric CO2 records from sites in the SIO air sampling network, in Trends: A Compendium of Data on Global Change, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tenn., 1998.*

[5] Berger, A., and M. F. Loutre, *Long-term variations in insolation and their effects on climate, the LLN experiments*, *Surveys in Geophysics*, 18 (2-3), 147-161, 1997.

[6] Berner, R. A., A. C. Lasaga, and R. M. Garrels, *The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years*, *American Journal of Science*, 283, 641-683, 1983.

[7] Prather, K, Noble CA. 1998. *Aerosol Time-of-Flight Mass Spectrometry: A New Method for Performing Real-Time Characterization of Aerosol Particles*. *Applied Occupational and Environmental Hygiene*. 13:439-443. [10.1080/1047322X.1998.10389569](https://doi.org/10.1080/1047322X.1998.10389569)

[8] Noble, C, Prather K. 1998. *Air pollution: the role of particles*. *Physics World*. 11:39-43.

[9] Houghton, J. T., L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, *Climate Change 1995, The Science of Climate Change*, Intergovernmental Panel on Climate Change, pp. 572, Cambridge University Press, Cambridge, GB, 1996.

[10] Ghil, M., and R. Vautard, *Interdecadal oscillations and the warming trend in global temperature time series*, *Science*, 199, 1065-1068, 1991.

[11] Hegerl, G. C., K. Hasselmann, U. Cubasch, J. F. B. Mitchell, E. Roeckner, R. Voss, and J. Waskewitz, *On multi-fingerprint detection and attribution of greenhouse gas and aerosol forced climatic change*, *Climate Dynamics*, 13, 613-634, 1997.

[12] Schlesinger, W. H., *Evidence from chronosequence studies for a low carbon-storage potential of soils*, *Nature*, 348, 232-234, 1990.

[13] Santer, B. D., K. E. Taylor, T. M. L. Wigley, J. E. Penner, P. D. Jones, and U. Cubasch, *Towards the detection and attribution of an anthropogenic effect on climate*, *Climate Dynamics*, 12, 79-100, 1995.

[14] Santer, B. D., T. M. L. Wigley, T. P. Barnett, and E. Anyamba, *Detection of climate change and attribution of causes, in Climate Change 1995: The Science of Climate Change*, edited by J. T. Houghton, L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, pp. 407-443, Cambridge University Press, Cambridge, UK, 1996.

[15] Hansen, J.E., and S. Lebedeff, 1987: *Global trends of measured surface air temperature*. *J. Geophys. Res.*, 92, 13345-13372, [doi:10.1029/JD092iD11p13345](https://doi.org/10.1029/JD092iD11p13345).

[16] Jones, Philip D; Jonsson, T; Wheeler, Dennis A (1997): *Monthly values of the North Atlantic Oscillation Index from 1821 to 2000*. PANGAEA.

[17] Mann, M. E., R. S. Bradley, and M. K. Hughes, *Global-scale temperature patterns and climate forcing over the past six centuries*, *Nature*, 392 (6678), 779-787, 1998.

[18] Hall, T.M., J.Y.N. Cho, M.C. Kelley, and W.K. Hocking,

1992: *A re-evaluation of the Stokes drift in the polar summer mesosphere*. *J. Geophys. Res.*, 97, 887-897, [doi:10.1029/91JD02835](https://doi.org/10.1029/91JD02835).

[19] Thompson, L. G., E. Mosely-Thompson, M. Davis, P. N. Lin, T. Yao, M. Dyrugerov, and J. Dai, "Recent Warming": *Ice core evidence from tropical ice cores with emphasis on central Asia*, *Global and Planetary Change*, 7, 145-156, 1993.

[20] Bayr, K. J., D. K. Hall, and W. M. Kovalick, *Observations on glaciers in the Eastern Austrian alps using satellite data*, *International Journal of Remote Sensing*, 15 (9), 1733-1742, 1994.

[21] Johannesson T., Sigurosson O. *Interpretation of glacier variations in Iceland 1930-1995*, National Energy Authority, Hydrological service, Grensasvegi 9, IS-108 Reykjavik.

[22] Hastenrath S., Greischar, LL. (1989): *Climatic Atlas of the Indian Ocean. Part III: Upper-Ocean Structure*. The University of Wisconsin, ISBN 0-299-12154-2.

[23] Bahr D. B., Dyrugerov M., and Meier F., *Sea-level rise from glaciers and ice caps: A lower bound*, *Geophysical research letters*, VOL. 36, L03501, [doi:10.1029/2008GL036309](https://doi.org/10.1029/2008GL036309), 2009

[24] Dahl-Jensen, D., K. Mosegaard, N. Gundestrup, G. D. Clow, S. J. Johnsen, A. W. Hansen, and N. Balling, *Past temperatures directly from the Greenland Ice Sheet*, *Science*, 282 (5387), 268-271, 1998.

[25] Berger, A., M. F. Loutre, and H. Gallee, *Sensitivity of the LLN climate model to the astronomical and CO2 forcings over the last 200 ky*, *Climate Dynamics*, 14 (9), 615-629, 1998

[26] Sundquist E.T. Broecker W.S., *The Carbon Cycle and Atmospheric CO2: Natural Variations Archean to Present*, Volume 32, :1 January 1985, Print ISBN:9780875900605 Online ISBN:9781118664322 DOI:10.1029/GM032

[27] Barnola, J. M., M. Anklin, J. Porcheron, D. Raynaud, J. Schwander, and B. Stauffer, *CO2 evolution during the last millennium as recorded by Antarctic and Greenland ice*, *Tellus*, 47B, 264-272, 1995.

[28] Delmotte, M., J. Chappellaz, D. Raynaud, E. Brook, and E. Tuttle, *Atmospheric methane changes during the last 400 kyr revealed by the Vostok ice core (abstract)*, *Supplement to Eos, Transactions, AGU*, 79 (45), F151-F152, 1998.

[29] Jouzel J, Barkov N. I., Barnola J. M., et. all, *Extending the Vostok ice-core record of palaeoclimate to the penultimate glacial period* Article Published: 29 July 1993, *Nature* volume 364, pages407-412.

[30] Berner, R. A., *The rise of plants and their effect on weathering and atmospheric CO2*, *Science*, 276, 544-546, 1997.

[31] Crowley, T. J., *The problem of paleo-analogs*, *Climatic Change*, 35 (1), 119-121, 1997.

[32] Tarasov, L., and W. R. Peltier, *Terminating the 100 kyr ice age cycle*, *J. Geophys. Res.- Atmos.*, 102 (D18), 21,665-21,693, 1997.

[33] Barron, E. J., *Explanations of the Tertiary global cooling trend*, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 50, 45-61, 1985.

[34] Boyle, E. A., *Chemical accumulation variations under the Peru current during the last 130,000 years*, *J. Geophys. Res.*, 88, 7667-7680, 1983.

[35] Sundquist, E. T., *Geologic analogs: Their value and limitations in carbon dioxide research*, in *The Changing Carbon Cycle, A Global Analysis*, edited by J. R. Trabalka, and D. E. Reiche, pp. 371-402, Springer-Verlag, New York, 1986.

[36] Sundquist, E. T., and W. S. Broecker, *The carbon cycle and atmospheric CO2: Natural variations Archean to present*, in *Geophysical Monograph*, pp. 627, American Geophysical Union, Washington, DC, 1985.

[37] Houghton J.T., Meira Filho L.G., Callander B.A., Harris N., Kattenberg A. and K. Maskell, *Climate Models – Evaluation*, *J Climate Change 1995, The Science of Climate Change*, Cambridge University.

[38] Krabill W., et al *Rapid Thinning of Parts of the Southern Greenland Ice Sheet*, April 1999 *Science* 83(5407):1522-4, DOI: 10.1126/science.283.5407.1522.

[39] Douglas, B. C., *Global sea level rise*, *J. Geophys. Res.*, 96 (C4), 6981-6992, 1991.

[40] Warrick, R. A., and J. Oerlemans, *Sea level rise, in Climate Change, The IPCC Assessment*, edited by J. T. Houghton, G. J. Jenkins, and J. J. Ephraums, Cambridge University Press, Cambridge, UK, 1990. ,

[41] Warrick, R. A., C. L. Provost, M. F. Meier, J. Oerlemans, and P. L. Woodworth, *Changes in sea level, Climate Change 1995: The Science of Climate Change*, 359-405, 1996.